

POSITION ARTICLE AND GUIDELINES

Open Access



# Early introduction is not enough: CSACI statement on the importance of ongoing regular ingestion as a means of food allergy prevention

Elissa M. Abrams<sup>1,2\*</sup> , Moshe Ben-Shoshan<sup>3</sup>, Jennifer L. P. Protudjer<sup>4,5,6,7,8</sup>, Elana Lavine<sup>9,10</sup> and Edmond S. Chan<sup>11</sup>

Both randomized controlled and observational studies demonstrate a reduction in the risk of food allergy with early introduction of common allergens, in particular peanut and egg [1–4]. While these studies vary in design, population, dose and allergen used, there is a strong and consistent trend that early introduction

has a role in the prevention of food allergy. As a result, guidelines clearly and consistently recommend early allergen ingestion (commonly operationalised as between ages 4–6 months), often in particular for higher risk populations, as a means of food allergy prevention [3, 5]. The current Canadian Society of Allergy and Clinical Immunology/Canadian Pediatric Society position statement recommends introduction of common allergens in high risk infants (with eczema or an immediate family history) at around 6, but not before 4 months of age when they are developmentally ready. In lower risk infants, introduction should be at around 6 months of age [5].

However, despite changes in guidance, recent epidemiologic evidence suggests that early introduction alone is not sufficient to reduce peanut allergy prevalence. In Australia, where uptake of peanut introduction in the first year of life increased more than threefold with adoption of early introduction guidelines, Soriano et al. reported observational data that peanut allergy prevalence had not significantly changed between 2007 and 2018 (3.1% to 2.6%, respectively; difference – 0.5% [95%CI – 1.4% to 0.4%];  $p=0.26$ ). This suggests that additional factors continue to play a role in the development of peanut and other food allergy (although with the guidelines only changing in 2016 studies may need longer to completely address this) [6].

While most Australian families were introducing peanut in infancy, Soriano reported that only ~30% of infants were eating peanut 2 or more times a week. A

\*Correspondence:

Elissa M. Abrams  
Elissa.abrams@gmail.com

<sup>1</sup> Department of Pediatrics, Section of Allergy and Clinical Immunology, University of Manitoba, FE125-685 William Avenue, Winnipeg, MB R3E 0Z2, Canada

<sup>2</sup> Department of Pediatrics, Division of Allergy and Immunology, University of British Columbia, 4480 Oak Street, Vancouver, BC V6H 3V4, Canada

<sup>3</sup> Division of Allergy Immunology and Dermatology, Department of Pediatrics, Montreal Children's Hospital, McGill University, Montreal, Canada

<sup>4</sup> Department of Pediatrics and Child Health, Max Rady College of Medicine, Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, MB, Canada

<sup>5</sup> Children's Hospital Research Institute of Manitoba, Winnipeg, MB, Canada

<sup>6</sup> Department of Food and Human Nutritional Sciences, Faculty of Agricultural and Food Sciences, University of Manitoba, Winnipeg, MB, Canada

<sup>7</sup> George and Fay Yee Centre for Healthcare Innovation, Winnipeg, MB, Canada

<sup>8</sup> Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

<sup>9</sup> Department of Pediatrics, University of Toronto, Toronto, Canada

<sup>10</sup> Department of Pediatrics, Queen's University, Humber River Hospital, Toronto, ON, Canada

<sup>11</sup> Division of Allergy, Department of Pediatrics, University of British Columbia, BC Children's Hospital, Vancouver, BC, Canada



substantial proportion were eating peanut less than once a week and some had even eaten peanut only once (a bite or taste). This suggests that a lack of regular ingestion may be a key reason for the lack of change in prevalence despite early introduction [7]. Although most allergy prevention guidelines, including the current CSACI/CPS guideline, recommend ongoing regular ingestion as a means of food allergy prevention [5, 8], this guidance is often not translated into practice, or individual recommendations to parents. Therefore, the goal of this statement is to emphasize that current best evidence supports the importance of regularity of infant allergen ingestion, operationalised as at least once weekly, as a means of food allergy prevention.

Immunologically, the premise of the “dual-allergen exposure hypothesis” is based on the need for oral tolerance to supersede cutaneous (or possibly respiratory) sensitization, and mechanistically this implies that consistent, regular oral ingestion is necessary to induce and maintain tolerance [9, 10]. Recent animal data suggests that regular peanut ingestion induces a regulatory T-cell population that express high levels of CTLA-4, which in turn suppresses T follicular helper cells and germinal center B-cells induced by environmental peanut exposure [11, 12]. All clinical studies on food allergy prevention have suggested both early introduction and ongoing regular ingestion to achieve optimal food allergy prevention. For example, in the Learning Early About Peanut (LEAP) study, infants in the early introduction group were introduced to peanut at age 4–11 months but also ate peanut at least 3 times a week (total of 6 g, or the equivalent of about 24 peanuts, per week) until age 5 years compared to strict avoidance [1]. Similarly, in the PETIT randomized controlled trial that compared early (age 6 months) vs delayed (age 12 months) egg introduction in infants with atopic dermatitis, the early introduction arm included ongoing feeding of egg at least daily [4]. Observational studies on early food introduction also support ongoing regular ingestion. For example, Katz et al. reported a large observational study in which delayed (after 14 days) and/or irregular (<1/day) cow’s milk ingestion significantly increased the risk of cow’s milk allergy (OR 19.3) compared to introduction in first 14 days of life [13]. In another case–control study of children with confirmed cow’s milk allergy (compared to nonatopic controls and children with egg allergy), irregularity of ingestion (<1/day) increased the risk of cow’s milk allergy [14]. A recent large prospective interventional study in which 1992 newborns were recruited shortly before birth to either exclusive breastfeeding or at least one meal of cow’s milk formula (with or without breastfeeding) daily for the first 2 months of life found a

significant reduction in cow’s milk allergy at 12 months (1.58% in the breastfed group compared to 0% in the other groups; relative risk 29.98,  $p < 0.001$ ) with regular cow’s milk formula ingestion. The vast majority of those who developed cow’s milk allergy while being breastfed were exposed to small amounts of cow’s milk formula during the first 2 months of life (prevalence of 0.7% in the per-protocol exclusively breastfed group versus 3.27% among infants exposed to small amounts of cow’s milk formula), resulting in the authors’ conclusion that early continuous exposure should be encouraged, while occasional exposure increases risk of IgE-mediated cow’s milk allergy and should be avoided [15]. Based on the cow’s milk literature, the CSACI/CPS position statement currently recommends that intermittent supplementation with intact cow’s milk formula should be avoided due to increased risk of cow’s milk allergy. While there is less evidence for allergens other than cow’s milk, egg and peanut, the mechanism of sensitization is felt to be the same for all common allergens and hence regularity of ingestion is likely to be beneficial for other common allergens including tree nuts, sesame, grains and seafood.

The randomized controlled trials and observational studies regarding early ingestion of allergens have focused on the infant population, but there are now emerging data in older children (in particular those at risk of allergy such as siblings of peanut allergic children) that irregularity of ingestion may also increase the risk of food allergy [16]. A dose-dependent relationship between frequency of peanut ingestion and reduced risk of peanut allergy was reported among siblings of peanut allergic children. These siblings had tolerated oral peanut challenges at baseline, and during a median 2.9 years of follow up, none (0%) of the siblings who ate peanut at least once a month had peanut allergy vs 3% who ate it less than once a month and 18% who completely avoided peanut [16].

Once an allergen is ingested, the exact frequency of allergen ingestion required to maintain tolerance is not yet clearly established. For example, in a recently published cluster-randomized trial, general population infants introduced to peanut, cow’s milk, wheat and egg from 3 months of age (but only eaten once a week; and less so after 6 months of age) had a significantly lower risk of food allergy than those introduced at standard age (no intervention) [17]. However, given the above updated evidence, a pragmatic recommendation is for the relevant allergen/s to be consumed multiple times per month (with a goal of at least once each week), integrated into a family’s diet. The amount of allergen required for ingestion also remains undetermined, although a secondary analysis of the Enquiring About Tolerance (EAT) randomized controlled trial suggested a dose

of about 2 g each of egg white and peanut protein per week (the equivalent of approximately one small boiled egg and 1.5 tsp peanut butter, respectively) is likely an amount that would be tolerated in infancy although further research is required and it is possible that smaller amounts may be effective as well [4, 18]. The long term duration of ingestion is unclear, but the LEAP-On study showed that when peanut was introduced in the 1 year of life and continued until 5 years of age, a subsequent 12 months of avoidance did not increase the prevalence of peanut allergy [19]. Based on this study, roughly 5 years of regular ingestion starting in infancy is likely a sufficient duration to induce long term tolerance.

In conclusion, our recommendations and considerations on frequency of ingestion are as follows (Fig. 1):

#### Recommendation:

1. Both early introduction and, once introduced, regular ingestion of age-appropriate amounts and textures of all common allergens multiple times per month (with a goal of at least once each week based on expert opinion) are very likely to be useful to establish and maintain tolerance.

#### Considerations:

Once introduced, current evidence suggests that a single exposure or occasional exposures could be detrimental and result in increased risk of sensitization and development of food allergy.

If an allergen is not a common component of the family's diet, and regular ingestion is not feasible for that family, avoidance may be preferable to intermittent ingestion although further research is required.

For newborn infants, exclusive breastfeeding is best, but if cow's milk formula is introduced it should be given regularly (e.g. one feeding per day to supplement breastfeeding) from that point forward rather than intermittently to prevent cow's milk allergy.

The ideal amount and frequency of regular ingestion remains unknown, but the above recommendation is based on a balance of evidence and practicality. In general an age-appropriate serving could be aimed for. A duration of 5 years appears to be enough to maintain tolerance to peanut, and other foods may require similar exposures.



**Fig. 1** The importance of ongoing regular ingestion as a means of food allergy prevention

#### Acknowledgements

The authors would like to acknowledge the CSACI Board Members P Begin, L Connors, A Ellis, K Hildebrand, H Kim, V Kim, M Latrous, D Mack, A O'Keefe, J Quirt, J Upton, T Vander Leek, H Vliagofitis for their assistance.

#### Author contributions

All authors contributed to the writing and/or editing of the manuscript.

#### Funding

Not applicable.

**Availability of data and materials**

Not applicable.

**Declarations****Ethical approval consent to participate.**

Not applicable.

**Competing interests**

EMA is an employee of Public Health Agency of Canada (PHAC); views expressed are her own and not those of PHAC. She is Section Head for Anaphylaxis/Food Allergy, and a Member of the Board, for the Canadian Society of Allergy and Clinical Immunology. JP is Section Head for Allied Health, and a Member of the Board, for the Canadian Society of Allergy and Clinical Immunology; sits on the steering committee for Canada's National Food Allergy Action Plan, and reports consultancy for Nutricia, Novartis and ALK-Abelló. MBS has received consultant and/or speaker fees from Medexus, Sanofi, Miravo, Stallergenes and Novartis. EL has been a member of advisory boards for Pfizer, Miravo, Medexus, Stallergenes and ALK; has provided consultant services to ALK. ESC has received research support from DBV Technologies; has been a member of advisory boards for Pfizer, Miravo, Medexus, Leo Pharma, Kaleo, DBV, AllerGenis, Sanofi Genzyme, Bausch Health, Avir Pharma, AstraZeneca, ALK; was an expert panel and coordinating committee member of the National Institute of Allergy and Infectious Diseases (NIAID)-sponsored Guidelines for Peanut Allergy Prevention; is co-lead of CPS/CSACI Food Allergy Prevention Position Statements; is on the Executive of the CSACI (Canadian Society of Allergy and Clinical Immunology); is on the Executive of the CPS (Canadian Paediatric Society) Allergy Section; and is a member of the healthcare advisory board for Food Allergy Canada.

Received: 20 March 2023 Accepted: 28 May 2023

Published online: 18 July 2023

**References**

- Du Toit G, Roberts G, Sayre PH, Bahnson HT, Radulovic S, Santos AF, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med*. 2015;372:803–13.
- Ierodiakonou D, Garcia-Larsen V, Logan A, Groome A, Cunha S, Chivinge J, et al. Timing of allergenic food introduction to the infant diet and risk of allergic or autoimmune disease: a systematic review and meta-analysis. *JAMA*. 2016;316:1181–92.
- Fleischer DM, Chan ES, Venter C, Spergel JM, Abrams EM, Stukus D, et al. A consensus approach to the primary prevention of food allergy through nutrition: guidance from the American academy of allergy, asthma, and immunology; American college of allergy, asthma, and immunology; and the Canadian society for allergy and clinica. *J allergy Clin Immunol Pract*. 2021;9:22–43.e4.
- Natsume O, Kabashima S, Nakazato J, Yamamoto-Hanada K, Narita M, Kondo M, et al. Two-step egg introduction for prevention of egg allergy in high-risk infants with eczema (PETIT): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2017;389:276–86.
- Abrams EM, Watson W, Vander Leek TK, Atkinson A, Primeau M-N, Francoeur M-J, et al. Dietary exposures and allergy prevention in high-risk infants. *Allergy Asthma Clin Immunol*. 2022;18:36.
- Soriano VX, Peters RL, Moreno-Betancur M, Ponsonby A-L, Gell G, Odoi A, et al. Association between earlier introduction of peanut and prevalence of peanut allergy in infants in Australia. *JAMA*. 2022;328:48–56.
- Soriano VX, Peters RL, Ponsonby A-L, Dharmage SC, Perrett KP, Field MJ, et al. Earlier ingestion of peanut after changes to infant feeding guidelines: the earlynuts study. *J Allergy Clin Immunol*. 2019;144:1327–1335.e5.
- Food Allergy Canada: Eat Early, Eat Often : <https://foodallergycanada.ca/wp-content/uploads/Eat-Early-Eat-Often.pdf>. Accessed 13 Mar 2023.
- Kulis MD, Smeekens JM, Immormino RM, Moran TP. The airway as a route of sensitization to peanut: an update to the dual allergen exposure hypothesis. *J Allergy Clin Immunol*. 2021;148:689–93.
- Lack G. Update on risk factors for food allergy. *J Allergy Clin Immunol*. 2012;129:1187–97.
- Moran TP, Kulis MD. A “LEAP” forward in understanding immune mechanisms of oral tolerance to peanut. *J Allergy Clin Immunol*. 2022;150:299–301.
- Krempski JW, Lama JK, Iijima K, Kobayashi T, Matsunaga M, Kita H. A mouse model of the LEAP study reveals a role for CTLA-4 in preventing peanut allergy induced by environmental peanut exposure. *J Allergy Clin Immunol*. 2022;150:425–439.e3.
- Katz Y, Rajuan N, Goldberg MR, Eisenberg E, Heyman E, Cohen A, et al. Early exposure to cow's milk protein is protective against IgE-mediated cow's milk protein allergy. *J Allergy Clin Immunol*. 2010;126:77–82.e1.
- Onizawa Y, Noguchi E, Okada M, Sumazaki R, Hayashi D. The association of the delayed introduction of cow's milk with IgE-mediated cow's milk allergies. *J Allergy Clin Immunol Pract*. 2016;4:481–488.e2.
- Lachover-Roth I, Cohen-Engler A, Furman Y, Shachar I, Rosman Y, Meir-Shafir K, et al. Early, continuing exposure to cow's milk formula and cow's milk allergy: the COMEET study, a single center, prospective interventional study. *Ann Allergy Asthma Immunol*. 2022;130:233.
- Paquin M, Paradis L, Graham F, Begin P, Des RA. Peanut consumption habits and incidence of new peanut allergy in a cohort of younger siblings of peanut-allergic children. *J allergy Clin Immunol Pract*. 2021;9:539–541.e1.
- Skjerven HO, Lie A, Vettukattil R, Rehbindler EM, LeBlanc M, Asaranoj A, et al. Early food intervention and skin emollients to prevent food allergy in young children (PreventADALL): a factorial, multicentre, cluster-randomised trial. *Lancet*. 2022;399:2398–411.
- Perkin MR, Logan K, Tseng A, Raji B, Ayis S, Peacock J, et al. Randomized trial of introduction of allergenic foods in breast-fed infants. *N Engl J Med*. 2016;374:1733–43.
- Du Toit G, Sayre PH, Roberts G, Sever ML, Lawson K, Bahnson HT, et al. Effect of avoidance on peanut allergy after early peanut consumption. *N Engl J Med*. 2016;374:1435–43.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

