ERRATUM Open Access



Erratum to: Two-weeks-sustained unresponsiveness by oral immunotherapy using microwave heated cow's milk for children with cow's milk allergy

Masaya Takahashi, Shoichiro Taniuchi^{*}, Kazuhiko Soejima, Yasuko Hatano, Sohsaku Yamanouchi and Kazuanri Kaneko

Erratum to: Allergy Asthma Clin Immunol (2016) 12:44 DOI 10.1186/s13223-016-0150-0

After publication of the original article [1], the author noticed that there are some error with Table 3 and Fig. 1.

The corrected Table 3 and Fig. 1 is given in this erratum. We apologize for any inconvenience this may cause.

Table 3 The rate of desensitization and 2-weeks-sustained unresponsiveness for cow's milk allergy in the oral immunotherapy group and the untreated group

| | OIT (N = 31) | | Untreated group $(N = 17)$ |
|---------------------|-----------------|--------------------------------------|------------------------------|
| | Desensitization | Two-weeks-sustained unresponsiveness | Pass OFC |
| At 1-year follow-up | 14/31 (45 %)*,# | 7/31 (21 %)** ^{, ##} | 0/17 (0 %)*,**, ^a |
| At 2-year follow-up | 18/30 (60 %)# | 14/30 (47 %)## | |
| At 3-year follow-up | 21/30 (70 %) | 16/30 (53 %) | |
| At 4-year follow-up | 17/20 (85 %) | 14/20 (70 %) | |

OIT oral immunotherapy, OFC open food challenge

^{*}Correspondence: taniuchi@hirakata.kmu.ac.jp Department of Pediatrics, Kansai Medical University, 2-5-1 Shin-machi, Hirakata, Osaka, Japan



^{*} P = 0.002 by Fisher's exact test

^{**} P = 0.036 by Fisher's exact test

 $^{^{\#}}$ P = 0.025 by Wilcoxon signed rank test

 $^{^{\#\#}}$ P = 0.008 by Wilcoxon signed rank test

^a Patients in untreated groups continued complete elimination of cow's milk and a year later were performed open food challenge using fresh cow's milk was performed




Fig. 1 Study enrollment and outcomes of *OIT* oral immunotherapy with *CM* cow's milk. (*Hash*) After the achievement of 2-weeks-sustained unresponsiveness, the patients are followed for 2–3 year and they are able to ingest CM and CM products freely without any adverse events. *OFC* open food challenge

The online version of the original article can be found under doi:10.1186/s13223-016-0150-0.

Reference

 Takahashi M, Taniuchi S, Soejima K, Hatano Y, Yamanouchi S, Kaneko K. Two-weeks-sustained unresponsiveness by oral immunotherapy using microwave heated cow's milk for children with cow's milk allergy. Allergy Asthma Clin Immunol. 2016;12:44. doi:10.1186/s13223-016-0150-0.

Published online: 31 October 2016