



POSTER PRESENTATION

Open Access

# Immunoregulatory role of secretory leukocyte protease inhibitor in allergic asthma

Rafael Marino<sup>1\*</sup>, Thusanth Thuraisingam<sup>2</sup>, Pierre Camateros<sup>1</sup>, Yong Zhong Xu<sup>1</sup>, Jennifer Henri<sup>3</sup>, Jingxuan Yang<sup>4</sup>, Guoan He<sup>5</sup>, Aihao Ding<sup>5</sup>, Danuta Radzioch<sup>6</sup>

From AllerGen NCE Inc.'s Fifth Annual Research Conference: Innovation from Cell to Society  
Québec City, QC, Canada. 7-9 February 2010

## Background

Asthma is a complex and multi-factorial inflammatory disease [1]. It is one of the most common chronic diseases among children and adolescents [2]. Secretory leukocyte protease inhibitor (SLPI) has shown higher levels in asthmatic patients and its function as an anti-inflammatory protein has been documented in respiratory diseases [3,4]. However, its role in the immunomodulation of the response during allergic asthma has not yet been fully elucidated. The aim of this study was to evaluate the role of SLPI in the development of phenotypes associated with allergic asthma, and the effect of resiquimod treatment on the SLPI and the possible mechanisms of action involved in the disease.

## Materials and methods

The importance of SLPI was assessed by evaluating airway resistance and inflammatory parameters in SLPI transgenic and knock-out mice using an ovalbumin (OVA)-induced model of acute allergic asthma and treatment with resiquimod.

## Results

Allergic SLPI transgenic mice showed a significant decrease in airway resistance compared to wild-type mice ( $6.3 \pm 1.1$  vs.  $8.0 \pm 2.1$  cm H<sub>2</sub>O × s/ml,  $p < 0.001$ ), the same effect was observed with inflammatory cell infiltration, eosinophil percentage ( $24 \pm 1.1\%$  vs.  $29 \pm 2.3\%$ ,  $p < 0.001$ ), goblet cells ( $6 \pm 1.4$  vs.  $36 \pm 4.0\%$ ,  $p < 0.001$ ) in the lungs and IgE levels ( $2014.1 \pm 309.2$  vs.  $4173.2 \pm 685.6$  ng/ml,  $p < 0.001$ ) in plasma. Allergic SLPI knock-out mice displayed significantly higher

values compared to wild-type mice. They include lung resistance ( $8.6 \pm 2.7$  vs.  $6.6 \pm 0.5$  cm H<sub>2</sub>O\*s/ml,  $p < 0.001$ ), inflammatory cell influx, eosinophils ( $36.0 \pm 2.7$  vs.  $29.0 \pm 1.5\%$ ,  $p < 0.001$ ), goblet cells ( $40 \pm 4.1$  vs.  $30 \pm 1.4\%$ ,  $p < 0.001$ ), cytokine levels in the lungs ( $p < 0.05$ ) and plasma IgE levels ( $3598 \pm 204.7$  vs.  $2763 \pm 220.3$  ng/ml,  $p < 0.001$ ). Expression of SLPI decreased inflammation in the lungs, plasma IgE levels, and lung resistance, whereas the ablation of SLPI has the opposite effect. Treatment with resiquimod improved airway resistance and inflammation of the lungs in SLPI knock-out and wild type, demonstrating that its effect is independent of the expression of SLPI.

## Conclusions

SLPI plays an immunoregulatory role in the respiratory tract by reducing the inflammatory process and by improving lung physiology in a murine model of acute allergic asthma.

## Acknowledgements

R. Marino and P. Camateros are both supported by a CIHR Doctoral Award. We acknowledge the histology expert assistance of Dr. Marie-Christine Guiot.

## Author details

<sup>1</sup>Department of Medicine, Montreal General Hospital, Research Institute, McGill University, Montreal, Quebec, Canada. <sup>2</sup>Faculty of Medicine, University of Montreal, Montreal, Quebec, Canada. <sup>3</sup>Montreal General Hospital, Research Institute, McGill University, Montreal, Quebec, Canada. <sup>4</sup>Institute of Cell Biology, Swiss Federal Institute of Technology Zürich, Zürich, Switzerland. <sup>5</sup>Department of Microbiology and Immunology, Weill Medical College of Cornell University, New York, New York, USA. <sup>6</sup>Faculty of Medicine, Department of Medicine and Department of Human Genetics, McGill University, Montreal, Quebec, Canada.

Published: 26 November 2010

## References

1. Myers TR: Use of heliox in children. *Respir Care* 2006, **51**:619-631.

\* Correspondence: rafael.marino@mail.mcgill.ca

<sup>1</sup>Department of Medicine, Montreal General Hospital, Research Institute, McGill University, Montreal, Quebec, Canada

Full list of author information is available at the end of the article

2. Blais L, Beauchesne MF, Levesque S: **Socioeconomic status and medication prescription patterns in pediatric asthma in Canada.** *J Adolesc Health* 2006, **38**:607-616.
3. Wang Z, Zheng T, Zhu Z, Homer RJ, Riese RJ, Chapman HA Jr, Shapiro SD, Elias JA: **Interferon gamma induction of pulmonary emphysema in the adult murine lung.** *J Exp Med* 2000, **192**:1587-1600.
4. Weldon S, McGarry N, Taggart CC, McElvaney NG: **The role of secretory leucoprotease inhibitor in the resolution of inflammatory responses.** *Biochem Soc Trans* 2007, **35**:273-276.

doi:10.1186/1710-1492-6-S3-P24

**Cite this article as:** Marino *et al.*: Immunoregulatory role of secretory leukocyte protease inhibitor in allergic asthma. *Allergy, Asthma & Clinical Immunology* 2010 **6**(Suppl 3):P24.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

